

Name:_____

NetID:_____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

(10 points) Suppose we have a function f defined by

$$\begin{aligned} f(0) &= f(1) = 3 \\ f(n) &= 5f(n-2) + d, \text{ for } n \geq 2 \end{aligned}$$

where d is a constant. Your partner has already figured out that

$$f(n) = 5^k f(n-2k) + \sum_{p=0}^{k-1} d5^p$$

Finish finding the closed form for $f(n)$ assuming that n is even. Show your work and simplify your answer. Recall the following useful closed form (for $r \neq 1$): $\sum_{k=0}^n r^k = \frac{r^{n+1} - 1}{r - 1}$

Solution:

To find the value of k at the base case, set $n - 2k = 0$. Then $n = 2k$, so $k = n/2$. Substituting this into the above, we get

$$\begin{aligned} f(n) &= 5^k f(n-2k) + \sum_{p=0}^{k-1} d5^p \\ &= 5^{n/2} \cdot 3 + d \sum_{p=0}^{n/2-1} 5^p \\ &= 3 \cdot 5^{n/2} + d \left(\frac{5^{n/2} - 1}{4} \right) \\ &= (3 + d/4)5^{n/2} - d/4 = (3 + d/4)(\sqrt{5})^n - d/4 \end{aligned}$$

Name: _____

NetID: _____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (8 points) Suppose we have a function f defined by

$$\begin{aligned} f(0) &= f(1) = 3 \\ f(n) &= 5f(n-2) + d, \text{ for } n \geq 2 \end{aligned}$$

where d is a constant. Express $f(n)$ in terms of $f(n-6)$ (where $n \geq 6$). Show your work and simplify your answer. You do **not** need to find a closed form for $f(n)$.

Solution:

$$\begin{aligned} f(n) &= 5f(n-2) + d \\ &= 5(5f(n-4) + d) + d \\ &= 5(5(5f(n-6) + d) + d) + d \\ &= 5^3 f(n-6) + (25 + 5 + 1)d \\ &= 5^3 f(n-6) + 31d \end{aligned}$$

2. (2 points) Check the (single) box that best characterizes each item.

$f(n) = n!$ can be defined recursively
by $f(0) = 1$, and $f(n) = nf(n-1)$
for all integers ...

$n \geq 0$ ☐

$n \geq 1$ ☒

$n \geq 2$ ☐

Name:_____

NetID:_____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (8 points) Suppose we have a function g defined (for n a power of 3) by

$$\begin{aligned} g(1) &= c \\ g(n) &= 3g(n/3) + n \text{ for } n \geq 3 \end{aligned}$$

Express $g(n)$ in terms of $g(n/3^3)$ (where $n \geq 27$). Show your work and simplify your answer. You do **not** need to find a closed form for $g(n)$.

Solution:

$$\begin{aligned} g(n) &= 3g(n/3) + n \\ &= 3(3g(n/9) + n/3) + n \\ &= 3(3(3g(n/27) + n/9) + n/3) + n \\ &= 27g(n/27) + n + n + n \\ &= 27g(n/27) + 3n \end{aligned}$$

2. (2 points) Suppose that $f : \mathbb{N} \rightarrow \mathbb{N}$ is such that $f(n) = n^2$. Give a recursive definition of f

Solution:

$$f(0) = 0, \text{ and } f(n+1) = f(n) + 2n + 1 \text{ for } n \geq 0.$$

You could also have used $f(n) = f(n-1) + 2n - 1$ for $n \geq 1$.

Name:_____

NetID:_____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

(10 points) Suppose we have a function g defined (for n a power of 4) by

$$\begin{aligned} g(1) &= c \\ g(n) &= 2g(n/4) + n \text{ for } n \geq 4 \end{aligned}$$

Your partner has already figured out that

$$g(n) = 2^k g(n/4^k) + n \sum_{p=0}^{k-1} \frac{1}{2^p}$$

Finish finding the closed form for $f(n)$ assuming that n is a power of 4. Show your work and simplify your answer. Recall that $\log_b n = (\log_a n)(\log_b a)$.

Solution: To find the value of k at the base case, set $n/4^k = 1$. Then $n = 4^k$, so $k = \log_4 n$. Notice also that $2^{\log_4 n} = 2^{\log_2 n \log_4 2} = n^{1/2} = \sqrt{n}$

Substituting this into the above, we get

$$\begin{aligned} g(n) &= 2^{\log_4 n} \cdot c + n \sum_{p=0}^{\log_4 n - 1} \frac{1}{2^p} \\ &= 2^{\log_4 n} \cdot c + n \left(2 - \frac{1}{2^{\log_4 n - 1}} \right) \\ &= c\sqrt{n} + n \left(2 - \frac{2}{\sqrt{n}} \right) \\ &= c\sqrt{n} + 2n - 2\sqrt{n} \\ &= 2n + (c - 2)\sqrt{n} \end{aligned}$$

Name: _____

NetID: _____

Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

(10 points) Suppose we have a function g defined (for n a power of 2) by

$$\begin{aligned} g(1) &= c \\ g(n) &= 4g(n/2) + n \text{ for } n \geq 2 \end{aligned}$$

Your partner has already figured out that

$$g(n) = 4^k g(n/2^k) + n \sum_{p=0}^{k-1} 2^p$$

Finish finding the closed form for $g(n)$ assuming that n is a power of 2. Show your work and simplify your answer. Recall that $\log_b n = (\log_a n)(\log_b a)$.

Solution: To find the value of k at the base case, set $n/2^k = 1$. Then $n = 2^k$, so $k = \log_2 n$. Notice also that $4^{\log_2 n} = 4^{\log_4 n \log_2 4} = n^2$

Substituting this into the above, we get

$$\begin{aligned} g(n) &= 4^{\log_2 n} \cdot c + n \sum_{p=0}^{\log_2 n - 1} 2^p \\ &= 4^{\log_2 n} \cdot c + n \sum_{p=0}^{\log_2 n - 1} 2^p \\ &= cn^2 + n(2^{\log_2 n} - 1) = cn^2 + n^2 - n = (c+1)n^2 - n \end{aligned}$$

Name: _____

NetID: _____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (8 points) Suppose we have a function g defined by

$$\begin{aligned} g(0) &= g(1) = c \\ g(n) &= kg(n-2) + n^2, \text{ for } n \geq 2 \end{aligned}$$

where k and c are constants. Express $g(n)$ in terms of $g(n-6)$ (where $n \geq 6$). Show your work and simplify your answer. You do **not** need to find a closed form for $g(n)$.

Solution:

$$\begin{aligned} g(n) &= kg(n-2) + n^2 \\ &= k(kg(n-4) + (n-2)^2) + n^2 \\ &= k(k(kg(n-6) + (n-4)^2) + (n-2)^2) + n^2 \\ &= k^3g(n-6) + k^2(n-4)^2 + k(n-2)^2 + n^2 \end{aligned}$$

2. (2 points) Check the (single) box that best characterizes each item.

The number of nodes in the
4-dimensional hypercube Q_4

4

☐

16

☒

32

☐

64

☐