

Name: _____

NetID: _____

Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (5 points) How many different 13-letter strings can be made by rearranging the letters in the word ‘‘massachusetts’’? Show your work.

2. (12 points) Check the (single) box that best characterizes each item.

If a function from \mathbb{R} to \mathbb{R} is strictly increasing, it must be one-to-one. true ☐ false ☐

$g : \mathbb{Z} \rightarrow \mathbb{Z}$
 $g(x) = 7 - \lfloor \frac{x}{3} \rfloor$ onto ☐ not onto ☐ not a function ☐

$g : (0, \frac{\pi}{2}) \rightarrow \mathbb{R}$
 $g(x) = \sin(x)$ one-to-one ☐ not one-to-one ☐ not a function ☐

We painted 12 mailboxes. There were 5 colors to choose from and each mailbox is painted with a single color. By the pigeonhole principle, every color appears on at least two mailboxes. true ☐ false ☐

$\exists y \in \mathbb{N}, \forall x \in \mathbb{N}, x = xy$ true ☐ false ☐

$(f \circ g)(x)$ $f(g(x))$ ☐ $g(f(x))$ ☐ neither ☐

Name: _____

NetID: _____

Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (5 points) Suppose that $|A| = p$, $|B| = q$, $|C| = n$. How many different functions are there from $A \times B$ to C ?

2. (12 points) Check the (single) box that best characterizes each item.

If a function from \mathbb{R} to \mathbb{R} is increasing,
it must be one-to-one.

true

☐

false

☐

$g : \mathbb{Z} \rightarrow \mathbb{R}$
 $g(x) = x + 2.137$

one-to-one

☐

not one-to-one

☐

not a function

☐

$g : \mathbb{Z} \rightarrow \mathbb{Z}$
 $g(x) = \lfloor x \rfloor$

onto

☐

not onto

☐

not a function

☐

Each ACM shirt has one of 6 trendy slogans. I bought
13 ACM shirts. Each slogan appears on at least two
shirts.

true

☐

false

☐

$\forall x \in \mathbb{Z}, \exists y \in \mathbb{Z}, x \neq y$ and $x + y = 0$

true

☐

false

☐

Suppose $f : A \rightarrow B$. For
all $x, y \in A$, if $f(x) = f(y)$,
then $x = y$.

onto

☐

one-to-one

☐

neither

☐

Name: _____

NetID: _____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (5 points) 10 men and 15 women showed up to this week's meeting of the UIUC Swing Dance Society. How many different ways can we form as many as possible into pairs, where each pair consists of one man and one woman?

2. (12 points) Check the (single) box that best characterizes each item.

If $f : A \rightarrow B$ is onto, then $|A| \geq |B|$ ☐ $|A| \leq |B|$ ☐ $|A| = |B|$ ☐

$f : \mathbb{Z} \rightarrow \mathbb{Z}$
 $f(x) = x + 3$ (x even),
 $f(x) = x - 21$ (x odd) onto ☐ not onto ☐ not a function ☐

$g : \mathbb{N} \rightarrow \mathbb{Z}$
 $g(x) = x^2$ one-to-one ☐ not one-to-one ☐ not a function ☐

Each ACM shirt has one of 6 trendy slogans. I bought 13 ACM shirts. At least three of these shirts must have the same slogan. true ☐ false ☐

$\forall x \in \mathbb{R}, \exists m, n \in \mathbb{Z}, x = \frac{m}{n}$ true ☐ false ☐

Suppose $f : A \rightarrow B$. For all $x \in A$, there is a $y \in B$,
 $f(x) = y$. onto ☐ one-to-one ☐ neither ☐

Name: _____

NetID: _____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (5 points) How many different 9-letter strings can be made by rearranging the letters in the word ‘‘silliness’’? Show your work.

2. (12 points) Check the (single) box that best characterizes each item.

A function is one-to-one if and only if each value in the co-domain has exactly one pre-image.

true ☐false ☐

$$f : \mathbb{Z} \rightarrow \mathbb{Z}$$

$$f(x) = x + 4 \text{ (} x \text{ even),}$$

$$f(x) = x - 21 \text{ (} x \text{ odd)}$$

one-to-one ☐not one-to-one ☐not a function ☐

$$f : \mathbb{N}^2 \rightarrow \mathbb{R}$$

$$f(p, q) = pq$$

onto ☐not onto ☐not a function ☐

Each ACM shirt has one of 6 trendy slogans. I bought 13 ACM shirts. There is a slogan that appears on at least two shirts.

true ☐false ☐

$$\exists y \in \mathbb{N}, \forall x \in \mathbb{Z}, x^2 = y$$

true ☐false ☐

Suppose $f : A \rightarrow B$. For all $x, y \in A$, if $x = y$, then $f(x) = f(y)$.

onto ☐one-to-one ☐neither ☐

Name: _____

NetID: _____

Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (5 points) Suppose that $|A| = 3$ and $|B| = 2$. How many onto functions are there from A to B ? Briefly justify or show work.

2. (12 points) Check the (single) box that best characterizes each item.

If a function is onto, then each value in the co-domain has at least one pre-image.

true

☐

false

☐

$$g : \mathbb{R} \rightarrow \mathbb{Z}$$

$$g(x) = \lfloor x \rfloor$$

one-to-one

☐

not one-to-one

☐

not a function

☐

$$g : \mathbb{Z} \rightarrow \mathbb{R}$$

$$g(x) = x - 0.314$$

onto

☐

not onto

☐

not a function

☐

Each dorm room is given an integer access code between 1 and 10 (inclusive). According to the pigeon-hole principle, if there are 21 dorm rooms, then there is some access code that is shared by at least two rooms.

true

☐

false

☐

$$\forall m, n \in \mathbb{Z}, \exists x \in \mathbb{Q}, x = \frac{m}{n}$$

true

☐

false

☐

Suppose $f : A \rightarrow B$. For all $y \in B$, there is an $x \in A$, $f(x) = y$.

onto

☐

one-to-one

☐

neither

☐