


1. Let  $C$  denote the curve pictured at right, with the orientation shown.

(a) For  $\mathbf{F}(x, y) = \langle xy, 0 \rangle$ , compute  $\int_C \mathbf{F} \cdot d\mathbf{r}$  directly. (3 points)



$$\int_C \mathbf{F} \cdot d\mathbf{r} =$$







(b) Check your answer to part (a) using Green's Theorem. (3 points)

2. For each function label its graph from among the options below: (2 points each)







(A)  $x^2 - y^2$

(B)  $\cos(xy)$

(C)  $e^{-(x^2+y^2)}$


☐

☐

☐

☐

☐

☐

3. (a) Each picture below depicts both (i) a constraint curve  $C$  defined by  $g(x, y) = 1$  for a function  $g(x, y)$ , and (ii) a level curve  $f(x, y) = M$  of a function  $f(x, y)$ . Mark the boxes of **all and only those pictures** for which  $M$  could be the maximum value of  $f(x, y)$  subject to the constraint  $g(x, y) = 1$ . [In every picture, you should assume that  $\nabla f$  is always nonzero.] **(2 points)**



- (b) Suppose a function  $f(x, y)$  attains its minimum value, subject to the constraint  $2x^2 + 2xy^2 + y^3 = 5$ , at  $(x, y) = (1, 1)$ . Assuming that  $\nabla f(1, 1) \neq \langle 0, 0 \rangle$ , find a nonzero vector  $\mathbf{v}$  parallel to  $\nabla f(1, 1)$ . **(3 points)**

$$\mathbf{v} = \left\langle \quad, \quad \right\rangle$$

4. Suppose  $f(x, y): \mathbb{R}^2 \rightarrow \mathbb{R}$  has the table of values and partial derivatives shown at right. For  $x(s, t) = s + 2t$  and  $y(s, t) = s^2 - t$ , let  $F(s, t) = f(x(s, t), y(s, t))$  be their composition with  $f$ . Compute  $\frac{\partial F}{\partial t}(2, 1)$ . **(3 points)**

$(x, y)$	$f(x, y)$	$\frac{\partial f}{\partial x}$	$\frac{\partial f}{\partial y}$
$(2, 1)$	0	7	6
$(2, -1)$	-12	7	-1
$(4, 3)$	7	3	1
$(5, 3)$	19	-8	5

$$\frac{\partial F}{\partial t}(2, 1) =$$







5. For each of the integrals

(A)  $\int_0^1 \int_0^1 \int_0^{1-x} f(x, y, z) \, dz \, dy \, dx$


(B)  $\int_0^1 \int_0^1 \int_0^y f(x, y, z) \, dx \, dy \, dz$

(C)  $\int_0^1 \int_x^1 \int_0^{y-x} f(x, y, z) \, dz \, dy \, dx$

label the solid corresponding to the region of integration below. (1 point each)




6. Compute the mass of solid region  $E$  shown at right if the mass density is  $\rho(x, y, z) = z$ . (4 points)



Mass =


7. (a) Let  $R$  be the region shown below right. Find a transformation  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  taking  $S = [0, 1] \times [0, 1]$  to  $R$ . (3 points)



$$T(u, v) = \left\langle \quad, \quad \right\rangle$$

- (b) Consider the transformation  $T(u, v) = (e^u - v, u + v)$  whose behavior is depicted below.

Compute  $\iint_R 3 \, dA$  via an integral over  $S$ . (3 points)



$$\iint_R 3 \, dA =$$

8. Let  $S$  be the surface in  $\mathbb{R}^3$  which is the boundary of the solid cube  $D = \{-1 \leq x \leq 1, -1 \leq y \leq 1, -1 \leq z \leq 1\}$ . For  $\mathbf{F}(x, y, z) = \langle yz^2 + e^z + x, ze^z + x + y, xe^x + xy + z \rangle$ , compute  $\iint_S \mathbf{F} \cdot \mathbf{n} \, dS$  by any valid method, where  $\mathbf{n}$  is the outward-pointing unit normal vector field. (4 points)


$$\iint_S \mathbf{F} \cdot \mathbf{n} \, dS =$$

9. Consider the region  $R$  below the surface  $z = 1 - x^2 - y^2$  and above the  $xy$ -plane. Compute the volume of  $R$ .  
(5 points)

Volume =

10. For each surface  $S$  in parts (a) and (b) give a parameterization  $\mathbf{r}: D \rightarrow S$ . Be sure to explicitly specify the domain  $D$  and call your parameters  $u$  and  $v$ .

- (a) The portion of the surface  $x = y^2$  shown at left. (2 points)



$D = \{$

$\mathbf{r}(u, v) = \langle$

- (b) The portion of the cylinder  $x^2 + z^2 = 1$  between the planes  $y = 0$  and  $y = 2$ . (3 points)

$D = \{$




$\mathbf{r}(u, v) = \langle$

- (c) Let  $M$  be the surface in part (b). Is the surface integral  $\iint_M y \, dS$ :  
 Circle your answer. (1 point)

negative   zero   positive

11. Let  $S$  be the surface parameterized by  $\mathbf{r}(u, v) = \langle u, uv, v \rangle$  for  $-1 \leq u \leq 1$  and  $-1 \leq v \leq 1$ .

(a) Mark the picture of  $S$  below. **(2 points)**



(b) Completely setup, but do not evaluate, the surface integral  $\iint_S x^2 \, dS$ . **(5 points)**




(c) Find the tangent plane to  $S$  at  $(0, 0, 0)$ . [You *must* show work that justifies your answer.] **(2 points)**

Equation:

$$\boxed{\phantom{0}}x + \boxed{\phantom{0}}y + \boxed{\phantom{0}}z = \boxed{\phantom{0}}$$

12. Consider the surface  $S$  parameterized by  $\mathbf{r}(u, v) = \langle \cos u, \sin u, v \rangle$  for  $0 \leq u \leq 2\pi$  and  $0 \leq v \leq 1$ .

(a) Mark the picture of  $S$  below. (2 points)



(b) Consider the vector field  $\mathbf{F} = \langle yz, -xz, 1 \rangle$  which has  $\text{curl} \mathbf{F} = \langle x, y, -2z \rangle$ . Directly evaluate  $\iint_S (\text{curl} \mathbf{F}) \cdot \mathbf{n} \, dS$  via the given parameterization, where  $\mathbf{n}$  is the outward normal vector field. (4 points)

$$\iint_S (\text{curl} \mathbf{F}) \cdot \mathbf{n} \, dS =$$

(c) Check your answer in (b) using Stokes' Theorem. (4 points)

13. Consider the function  $f(x, y)$  on the rectangle  $D = \{0 \leq x \leq 4 \text{ and } 0 \leq y \leq 2\}$  whose graph is shown below right. For each part, circle the best answer. (1 point each)

(a) At the point  $P = (1, 0.5)$  is  $\frac{\partial f}{\partial y}$ :

negative zero positive

(b) At  $P$  is  $\frac{\partial^2 f}{\partial x^2}$ :

negative zero positive

(c) How many critical points does  $f$  have in the interior of  $D$ ?

0 1 2 3 4

(d) The integral  $\iint_D f(x, y) dA$  is:

negative zero positive






(e) For the curve  $C$  shown, the line integral  $\int_C \nabla f \cdot d\mathbf{r}$  is:

-3 -1.5 0 1.5 3

(f) The line integral  $\int_C f ds$  is:

negative zero positive

(g) Mark the plot of the vector field  $\nabla f$ .








14. For each problem, circle the best answer. (1 point each)

(a) Consider the vector field  $\mathbf{F} = \langle 1, x, -z \rangle$ . The vector field  $\mathbf{F}$  is:

conservative      not conservative

(b) Mark the plot of  $\mathbf{F}$  on the region where each of  $x, y, z$  is in  $[0, 1]$ :




(c) For the leftmost vector field in part (b) is the divergence:

negative      zero      positive

Let  $S$  and  $H$  be the surfaces at right; the boundary of  $S$  is the unit circle in the  $xy$ -plane, and  $H$  has no boundary. Let  $\mathbf{G} = \langle x, y, z \rangle$ .

(d) The flux  $\iint_H \mathbf{G} \cdot \mathbf{m} \, dS$  is:

negative      zero      positive



(e) The flux  $\iint_S \mathbf{G} \cdot \mathbf{n} \, dS$  is:

negative      zero      positive

(f) The flux  $\iint_S (\text{curl } \mathbf{G}) \cdot \mathbf{n} \, dS$  is:

negative      zero      positive